

Smoking and bone health

© 2024 National Centre for Smoking Cessation and Training (NCSCT)

Author: Lion Shahab

Editor: Andy McEwen

Review of literature (October 2024): Sophia Papadakis

Executive summary

Poor bone health has a devastating impact in the UK, both in terms of disease morbidity and mortality as well as financial costs. Smoking has long been acknowledged to be a risk factor for poor bone health as it affects the metabolism of hormones, body weight, vitamin D levels, calcium absorption, blood circulation and increases oxidative stress thus disrupting healthy bone resorption and formation, leading to osteoporosis. Consequently, people who smoke have a 25% increase in fracture risk and are nearly twice as likely to experience hip fractures. Smoking also delays bone healing following operations to repair fractures. Stopping smoking has been shown to partially reverse the risk of suffering fractures, and smoking cessation is therefore advised in national guidelines for the prevention and treatment of osteoporosis.

Key points

1. Bone health – the scale of the problem


- The main disease associated with poor bone health, osteoporosis, is diagnosed in over 3 million people in UK, or about 5.2% of the UK population. Prevalence increases steeply with age: one in two women and one in five men over the age of 50 are affected by the disease.^{1,2}
- Osteoporosis is characterised by low bone mineral density (BMD) and deterioration of bone tissue, which leads to progressive bone fragility and causes over 530,000 fractures per year in the UK, primarily of the spine, hip, wrist, humerus and pelvis.^{1,2}
- Fractures greatly affect morbidity and mortality; up to a third die during the first year following a hip fracture, another third require nursing home placements and less than a third will regain a normal level of physical function.³⁻⁷ Fractures also result in increased back pain, height loss and physical disability.³⁻⁵
- The costs of fractures are high, currently totalling more than £4.4 billion per year in the UK, including £2 billion for treating hip fractures alone.¹⁻³

2. Smoking and bone health

2.1 Primary effects

- Bone health is primarily determined by peak bone mass achieved (usually around 30 years of age) and the rate of bone loss in the succeeding years.⁸ While the former is largely dependent on untreatable factors such as genetics, the later is not only determined by non-modifiable causes like age but also by modifiable risk factors such as physical inactivity.⁹
- Among treatable causes of osteoporosis, smoking has long been established as a contributing risk factor¹⁰ as it affects the balance of the naturally occurring processes of bone resorption and bone formation, resulting in low BMD as the amount resorbed is not fully replaced.^{8,11–13}
- Smoking is thought to cause low bone density through various pathways (see Figure 1): (1) Smoking has been linked to changes in hormone household, leading to a decrease in parathyroid hormone (thus reducing calcium absorption) and oestrogen levels as well as to an increase in the level of cortisol and adrenal androgens, changes that have been linked to an increased risk of osteoporosis¹⁴; (2) Smoking reduces body mass, which is postulated to provide an osteogenic stimulus and is linked to higher BMD¹⁵; (3) Smoking reduces the level of Vitamin D in the body, which is required for good bone health¹⁶; (4) Smoking increases free radicals and oxidative stress which affects bone resorption¹⁷; (5) People who smoke are more likely to suffer from peripheral vascular disease, reducing blood supply to the bones¹⁸; (6) As people who smoke are weaker, have poorer balance and impaired neuromuscular performance, smoking may also increase the risk of falls¹⁹; (7) Finally, there may also exist direct toxic effects of many of the constituents in tobacco smoke on bone cells.²⁰
- Meta-analyses have attempted to estimate the effect of smoking on bone health. While estimates vary (see Figure 2), there is a significant effect of smoking on overall fracture risk – in particular for the hip, spine and heel bone.^{18,21–24} Overall, risk of any fracture is increased by about 25% in people currently smoking and for hip fractures risk is increased between 40–84% and there is an increase in risk to over 100% in those over 85 years of age.^{18,21–24} Meta-analysis data has also found smoking is associated with increased mortality in patients with hip fracture.²⁵

Figure 1. Potential mechanisms of increased fracture risk among smokers.
Adapted from Wong et al, 2007.¹⁰



SHBG – sex hormone-binding globulin


2.2 Secondary effects

- As a consequence of osteoporosis, those affected often require surgery to deal with the complications of fractures. There is convincing evidence to suggest that smoking is linked to impaired bone healing.^{26–29} This is most likely due to its potential impact on cellular differentiation and compromised microcirculation, both required for fracture repair.²⁹

3. Smoking cessation and bone health

- While life-time risk of fractures is increased in people who used to smoke compared with people who have never smoked, this risk is lower than among people who currently smoke (Figure 2). Several studies have shown that BMD is intermediate between people who currently smoke and people who have never smoked, suggesting that the effects of smoking may be partially reversible^{11,30–31} and there is a dose-response relationship for the amount in pack-years smoked and fracture risk.³²
- Longitudinal studies have shown that smoking cessation reduces bone loss^{21,23,33} and smoking cessation is therefore recommended in osteoporosis guidelines.^{3,34}

Figure 2. Fracture risk associated with current and past smoking.



A) Data from Vestergaard and Mosekilde, 2003¹⁸

B) Data from Kanis et al, 2005²¹

References

- Svedbom A, Helmlund E, Ivergård M, Compston J, Cooper C, Stenmark J, et al. Osteoporosis in the European Union: a compendium of country specific reports. *Arch Osteoporos* [Internet]. 2013;8(1–2). Available from: www.ncbi.nlm.nih.gov/pmc/articles/PMC3880492/
- International Osteoporosis Foundation. Epidemiology, Burden, and Treatment of Osteoporosis in the United Kingdom. International Osteoporosis Foundation:2021. Available from: <https://www.osteoporosis.foundation/sites/IOFbonehealth/files/scope-2021/UK%20report.pdf>
- Royal Osteoporosis Society. Effective secondary prevention of fragility fractures: clinical standards for fracture liaison services. Royal Osteoporosis Society; 2019. Available from: <https://theros.org.uk/media/teubz33w/ros-clinical-standards-for-fracture-liaison-services-august-2019.pdf>
- Public Health England (PHE). Falls and fracture consensus statement; supporting commissioning for prevention. PHE; 2017. Available from: https://assets.publishing.service.gov.uk/media/5a7f4316ed915d74e6229625/falls_and_fractures_consensus_statement.pdf
- Royal College of Physicians. 15 years of quality improvement: The 2023 National Hip Fracture Database report on 2022 RCP:London:2023. Available from: <https://www.nhfd.co.uk/2023report>
- Bliuc D, Nguyen ND, Milch VE, Nguyen TV, Eisman JA, Center JR. Mortality risk associated with low-trauma osteoporotic fracture and subsequent fracture in men and women. *JAMA* 2009;301:513–521
- Neuberger J, Currie C, Wakeman R, Tsang C, Plant F, De Stavola B, et al. The Impact of the National Clinician-led Audit Initiative on Care and Mortality after Hip Fracture in England. *Medical Care* 2015;53(8):686–91.
- Office of the Surgeon General (US). Bone Health and Osteoporosis: A Report of the Surgeon General. Rockville (MD): Office of the Surgeon General (US); 2004. 2. The Basics of Bone in Health and Disease. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK45504/>
- Cummings SR. Treatable and untreatable risk factors for hip fracture. *Bone* 1996;18(3 Suppl):165S–167S.
- Wong PK, Christie JJ, Wark JD. The effects of smoking on bone health. *Clin Sci (Lond)* 2007;113(5):233–241.
- Szulc P, Garnero P, Claustrat B, Marchand F, Duboeuf F, Delmas PD. Increased bone resorption in moderate smokers with low body weight: the Minos study. *J Clin Endocrinol Metab* 2002;87(2):666–674.
- Hernigou J, Schuind F. Tobacco and bone fractures: A review of the facts and issues that every orthopaedic surgeon should know. *Bone Joint Res.* 2019 Jul 5;8(6):255–265.
- Pompe E, Bartstra J, Verhaar HJ, de Koning HJ, van der Aalst CM, Oudkerk M, Vliethehart R, Lammers JJ, de Jong PA, Mohamed Hoesein FAA. Bone density loss on computed tomography at 3-year follow-up in current compared to former male smokers. *Eur J Radiol.* 2017 Apr;89:177–181. doi: 10.1016/j.ejrad.2017.02.011. Epub 2017 Feb 9. PMID: 28267536.
- Kapoor D, Jones TH. Smoking and hormones in health and endocrine disorders. *Eur J Endocrinol* 2005; 152(4):491–499.
- Daniel M, Martin AD, Drinkwater DT. Cigarette smoking, steroid hormones, and bone mineral density in young women. *Calcif Tissue Int* 1992;50(4):300–305.
- Brot C, Jorgensen NR, Sorensen OH. The influence of smoking on vitamin D status and calcium metabolism. *Eur J Clin Nutr* 1999;53(12):920–926.
- Duthie GG, Arthur JR, James WP. Effects of smoking and vitamin E on blood antioxidant status. *Am J Clin Nutr* 1991;53(4 Suppl):1061S–1063S.
- Vestergaard P, Mosekilde L. Fracture risk associated with smoking: a meta-analysis. *J Intern Med* 2003;254(6):572–583.
- Nelson HD, Nevitt MC, Scott JC, Stone KL, Cummings SR. Smoking, alcohol, and neuromuscular and physical function of older women. Study of Osteoporotic Fractures Research Group. *JAMA* 1994;272(23):1825–1831.
- Broulik PD, Jarab J. The effect of chronic nicotine administration on bone mineral content in mice. *Horm Metab Res* 1993;25(4):219–221.
- Kanis JA, Johnell O, Oden A, Johansson H, De Laet C, Eisman JA et al. Smoking and fracture risk: a meta-analysis. *Osteoporos Int* 2005;16(2):155–162.
- Ward KD, Klesges RC. A meta-analysis of the effects of cigarette smoking on bone mineral density. *Calcif Tissue Int* 2001; 68(5):259–583.
- Wu ZJ, Zhao P, Liu B, Yuan ZC. Effect of Cigarette Smoking on Risk of Hip Fracture in Men: A Meta-Analysis of 14 Prospective Cohort Studies. *PLoS One.* 2016 Dec 30;11(12):e0168990.
- Law MR, Hackshaw AK. A meta-analysis of cigarette smoking, bone mineral density and risk of hip fracture: recognition of a major effect. *BMJ* 1997;315(7112):841–846.
- Zhang N, Liu YJ, Yang C, Zeng P, Gong T, Tao L, Li XA. Association between cigarette smoking and mortality in patients with hip fracture: A systematic review and meta-analysis. *Tob Induc Dis.* 2022;20:110.
- Haverstock BD, Mandracchia VJ. Cigarette smoking and bone healing: implications in foot and ankle surgery. *J Foot Ankle Surg* 1998;37(1):69–74.
- Patel RA, Wilson RF, Patel PA, Palmer RM. The effect of smoking on bone healing: A systematic review. *Bone Joint Res.* 2013 Jun 14;2(6):102–11.
- Xu B, Anderson DB, Park ES, Chen L, Lee JH. The influence of smoking and alcohol on bone healing: Systematic review and meta-analysis of non-pathological fractures. *EClinicalMedicine.* 2021 Oct 31;42:101179.
- Kwong FN, Harris MB. Recent developments in the biology of fracture repair. *J Am Acad Orthop Surg* 2008;16(11):619–625.
- Trevisan C, Alessi A, Girotti G, Zanforlini BM, Bertocco A, Mazzochin M, et al. The Impact of Smoking on Bone Metabolism, Bone Mineral Density and Vertebral Fractures in Postmenopausal Women. *J Clin Densitom.* 2020;23(3):381–389.
- Nguyen TV, Kelly PJ, Sambrook PN, Gilbert C, Pocock NA, Eisman JA. Lifestyle factors and bone density in the elderly: implications for osteoporosis prevention. *J Bone Miner Res* 1994;9(9):1339–1346.
- Valtola A, Honkanen R, Kroger H, Tuppurainen M, Saarikoski S, Alhava E. Lifestyle and other factors predict ankle fractures in perimenopausal women: a population-based prospective cohort study. *Bone* 2002;30(1):238–242.
- Hollenbach KA, Barrett-Connor E, Edelstein SL, Holbrook T. Cigarette smoking and bone mineral density in older men and women. *Am J Public Health* 1993;83(9):1265–1270.
- Gregson CL, Armstrong DJ, Bowden J, Cooper C, Edwards J, Gittoes NJL, et al. UK clinical guideline for the prevention and treatment of osteoporosis. *Arch Osteoporos.* 2022 Apr 5;17(1):58. doi: 10.1007/s11657-022-01061-5.