

The Clinical Case for providing stop smoking support to Hospitalised Patients

Why intervene in secondary care?

1. Hospital patients are more receptive to 'Very Brief Advice' (VBA) and an offer of support to stop smoking, as they are often experiencing a period of heightened motivation
2. Giving VBA to a hospital patient (the '3 A's': Ask, Advise, Act) can also encourage compliance to the smokefree hospital policy, and highlight any need for withdrawal management. Providing Nicotine Replacement Therapy (NRT) to a patient during a period of forced abstinence, will ease nicotine withdrawal symptoms
3. Stopping smoking can lead to significant health benefits, and reduce post-operative complications and improve recovery time

What is the aim of this 'clinical case' document?

The aim of this document is to provide clinical support for hospital staff in terms of supporting patients to stop smoking, even if this is just for a period of forced abstinence whilst in hospital. Being in hospital provides an opportune moment to intervene and provide both brief advice and support to stop smoking; including making a referral on to local stop smoking support. There are many benefits for a patient if they have temporary abstinence from smoking, including a shorter time for recovery and this can often stimulate a full attempt to stop smoking.

What is the relationship between smoking and hospitalisation?

Smoking is a major contributor to hospitalisation.¹ In one large population-based study, increased short-term rates of hospitalisation in a young healthy population were attributable to smoking.² It has been estimated that eliminating smoking would reduce annual rates of all-cause hospitalisation among older adults by 9%.³

Effects of hospitalisation on stopping smoking

- The experience of hospitalisation itself led to a substantial number of patients stopping smoking long-term.^{4,5}
- Many patients stop smoking during hospital admissions and those who do not stop completely, tend to decrease their use of tobacco.⁶

The Clinical Case for providing stop smoking support to Hospitalised Patients

What are the health benefits of stopping smoking for hospitalised patients?

Stopping smoking will not only benefit a patient's long term health by reducing the risk of developing smoking related disease,⁷ but smoking abstinence may also help a patient to recover quicker by eliminating the acute effects of smoking on the body. There is an evidenced benefit of stopping smoking in terms of hospitalisation outcomes and general outcomes (see below).

Main acute effects of smoking on the body (estimated time of recovery, if known)

- Increase in sympathetic tone leading to an increase in blood pressure, heart rate and peripheral vasoconstriction leading to an increased demand for oxygen and cardiac function.⁸
(24 – 48 hours)
- Formation of carboxyhaemoglobin leading to a reduction in oxygen delivery to the tissues.⁹
(8 – 24 hours)
- Formation of carboxymyoglobin leading to a reduction in oxygen storage in the muscles.¹⁰
(8 – 24 hours)
- Increase in red blood cell production, which leads to an increase in blood viscosity, a decrease in tissue perfusion, a decrease in oxygen delivery to the tissues and potentiation of thrombotic process.^{11,8}
- Hypersecretion of mucus, narrowing of the small airways, decrease in ciliary function and change in mucus rheology leading to a decrease in mucociliary transport.^{11,8} **(12 – 72 hours)**
- Changes in functioning of a range of immune cells (pro- and anti-inflammatory cytokines, white blood cells, immunoglobulins) which lead to decreased immunity and are associated with atherosclerosis.^{11,8} **(1 week – 2 months)**
- Induction of hepatic enzymes which increases drug metabolism through both pharmacokinetic and pharmacodynamic mechanisms.¹² **(6 – 8 weeks)**

The Clinical Case for providing stop smoking support to Hospitalised Patients

Effects of stopping smoking on hospitalisation

- Risk of hospitalisation decreases to the level of non-smokers in 1 to 3 years after quitting.^{13,1}
- Stopping smoking decreases the risk of postoperative complications and reduces wound healing complications. Stopping smoking for a longer duration compared to shorter periods has an average reduction in total complications of 20%.¹⁴

General health benefits of stopping smoking¹⁵

- Within 20 minutes blood pressure drops to the level it was before the last cigarette.
- Within 8 hours carbon monoxide levels in the blood return to normal.
- Within 24 hours the chance of a heart attack decreases.
- Within 2 weeks to 3 months circulation improves and lung function increases.
- Within 1 to 9 months lungs regain normal ciliary function, reducing infection risk.
- By 10 years the risk of lung cancer is approximately half of a smoker. The risk of cancers of the mouth, throat, bladder, kidney and pancreas also decrease.

Stop smoking support is effective

Providing stop smoking support has been proven effective for hospitalised patients, regardless of reason for admission.¹⁶

Effective methods typically include a combination of medications, replacing unhealthy habits with healthy habits and behavioural support. Evidence indicates that NRT, bupropion and varenicline are all effective treatments for stopping smoking both short and long-term.¹⁷

A 2008 Cochrane review demonstrated the positive impact of implementing stop smoking services for hospital patients.¹⁶

The Clinical Case for providing stop smoking support to Hospitalised Patients

Providing 'Very Brief Advice' to hospital patients: the '3 A's'

Providing a stop smoking intervention to a hospital patient is proven to be effective regardless of the reason for admission.¹⁸ Offering VBA is the single most cost effective and clinically proven preventative action a healthcare professional can take¹⁹ and it is important to keep giving advice at every opportunity, as smokers may take several attempts to stop smoking successfully.²⁰ In addition, by referring a patient to a local stop smoking service, they are four times more likely to stop smoking.²¹

Research shows that 95% of patients expect to be asked about smoking and a short intervention can make all the difference.^{22,23} The '3 A's' 30 second approach to giving 'very brief advice' are as follows:

ASK and record smoking status

ADVISE the patient of the personal health benefits of stopping smoking

ACT on the patient's response

- prescribe NRT for patients in withdrawal
- monitor withdrawal and adjust pharmacotherapy accordingly
- refer to local stop smoking service

How was this information sheet put together?

This information is a summary of the current scientific evidence on the association between cigarette smoking and hospitalisation. Studies were found by searching MEDLINE and EMBASE using combined exploded subject headings of 'hospitalisation' and 'tobacco use cessation' from 01/1945 – 07/2011. Evidence has been included in this summary from large population based studies, cohort studies, randomised controlled trials and reviews only.

The Clinical Case for providing stop smoking support to Hospitalised Patients

References

- English, D. R., H. T. Vu, and M. W. Knuiman. 2002. "The impact of smoking on use of hospital services: the Busselton study." *Aust N Z J Public Health* 26(3):225–30.
- Robbins, A. S., V. P. Fonseca, S. Y. Chao, G. A. Coil, N. S. Bell, and P. J. Amoroso. 2000. "Short term effects of cigarette smoking on hospitalisation and associated lost workdays in a young healthy population." *Tob Control* 9(4):389–96.
- Russell, L. B., S. M. Teutsch, R. Kumar, A. Dey, and E. Milan. 2001. "Preventable smoking and exercise-related hospital admissions. A model based on the NHEFS." *Am J Prev Med* 20(1):26–34.
- Lando, H., D. Hennrikus, M. McCarty, and J. Vessey. 2003. "Predictors of quitting in hospitalized smokers." *Nicotine Tob Res* 5(2):215–22.
- Ong, K. C., G. N. Cheong, L. Prabhakaran, and A. Earnest. 2005. "Predictors of success in smoking cessation among hospitalized patients." *Respirology* 10(1):63–9.
- Duffy, S. A., R. L. Scholten, and C. A. Karvonen-Gutierrez. 2010. "The relation of tobacco use during hospitalization to post-discharge smoking cessation among US veterans." *Prev Med* 50(5–6):285–7.
- Doll, R., R. Peto, J. Boreham, and I. Sutherland. 2004. "Mortality in relation to smoking: 50 years' observations on male British doctors." *BMJ* 328(7455):1519.
- Warner, D. O. 2006. "Perioperative abstinence from cigarettes: physiologic and clinical consequences." *Anesthesiology* 104(2):356–67.
- Rietbrock, N., S. Kunkel, W. Wörner, and P. Eyer. 1992. "Oxygen-dissociation kinetics in the blood of smokers and non-smokers: interaction between oxygen and carbon monoxide at the hemoglobin molecule." *Naunyn Schmiedebergs Arch Pharmacol* 345(1):123–8.
- Akrawi, W., and J. L. Benumof. 1997. "A pathophysiological basis for informed preoperative smoking cessation counseling." *J Cardiothorac Vasc Anesth* 11(5):629–40.
- Moller, H., and H. Tonnesen. 1997. "Alcohol drinking, social class and cancer." *IARC Sci Publ* (138):251–63.
- Zevin, S., and N. L. Benowitz. 1999. "Drug interactions with tobacco smoking. An update." *Clin Pharmacokinet* 36(6):425–38.
- Baumeister, S. E., A. Schumann, C. Meyer, U. John, H. Volzke, and D. Alte. 2007. "Effects of smoking cessation on health care use: is elevated risk of hospitalization among former smokers attributable to smoking-related morbidity?" *Drug Alcohol Depend* 88(2–3):197–203.
- Mills, E., O. Eyawo, I. Lockhart, S. Kelly, P. Wu, and J. O. Ebbert. 2011. "Smoking cessation reduces postoperative complications: a systematic review and meta-analysis." *Am J Med* 124(2):144–54 e8.
- Shah, R. S., and J. W. Cole. 2010. "Smoking and stroke: the more you smoke the more you stroke." *Expert Rev Cardiovasc Ther* 8(7):917–32.
- Rigotti, N. A., M. R. Munafo, and L. F. Stead. 2008. "Smoking cessation interventions for hospitalized smokers: a systematic review." *Arch Intern Med* 168(18):1950–60.
- Wu, P., K. Wilson, P. Dimoulas, and E. J. Mills. 2006. "Effectiveness of smoking cessation therapies: a systematic review and meta-analysis." *BMC Public Health* 6:300.
- Rigotti N, Munafo MR, Stead LF. Interventions for smoking cessation in hospitalised patients. *Cochrane Database of Systematic Reviews* 2007; Issue3.Art.No.:CD001837.DOI:10.1002/14651858.CD001837.pub2.
- Anczakj, Nogler (2003) . Tobacco cessation in primary care: maximizing intervention strategies. *Clinical Medicine & Research* 2003; 1: 201–216
- Fu S, Partin M, Snyder A, An LC, Nelson DB, Clothier B, Nugent S, Willenbring ML, Joseph AM. (2006) Promoting repeat tobacco dependence treatment: are relapsed smokers interested? *American Journal of managed Care* 2006; 12 235–243
- Smoking Toolkit Study (2001) Available at: <http://www.smokinginengland.info/>
- Slama KJ, Redman S, Cockburn J, Sanson-Fisher R. Community views about the role of general practitioners in disease prevention. *Family Practice* 1989; 6: 203–209.
- Department of Health 2009. "Stop Smoking Interventions in secondary care." in www.dh.gov.uk/publications.